

Gaming the Quantum

Faisal Shah Khan, Simon J.D. Phoenix
Khalifa University, Abu Dhabi, UAE

February 10, 2012

Abstract

In the time since a merger of quantum mechanics and game theory was proposed formally in 1999, the two distinct perspectives apparent in this merger of applying quantum mechanics to game theory, referred to henceforth as the theory of “quantized games”, and of applying game theory to quantum mechanics, referred to henceforth as “gaming the quantum”, have become synonymous under the single ill-defined term “quantum game”. Here, these two perspectives are delineated and a game-theoretically proper description of what makes a multi-player, non-cooperative game quantum mechanical, is given. Within the context of this description, finding a Nash equilibrium in a quantum game is shown to be equivalent to finding a solution to a simultaneous best approximation problem in the state space of quantum objects, thus setting up a framework for a game theory inspired study of “equilibrium” behavior of quantum physical systems such as those utilized in quantum information processing and computation.

1 Multi-player, Non-cooperative Games

Multi-player, non-cooperative game theory can be described as the science of making optimal choices under given constraints. To this end, introduce a set O of *outcomes*, a finite number, say n , of individuals called *players* with non-identical preferences over these outcomes, and assume that the players interact with each other within the constraints of these outcomes. Call this interaction between players a *game* and define it to be a function G with range equal to the set O and domain equal to the Cartesian product $S_1 \times S_2 \times \cdots \times S_n$ with S_i defined to be the set of the i^{th} players *pure strategies*. Call an element of the set $S_1 \times S_2 \times \cdots \times S_n$ a *play of the game* or a *strategy profile*. In symbols, a game G is a function

$$G : S_1 \times S_2 \times \cdots \times S_n \rightarrow O.$$

It is also assumed that the players make strategic choices independently. Note that a game is distinguished from an ordinary function by the notion of non-identical preferences, one per player, defined over the elements of its range or the outcomes.

Given a game G as above, assume further that the each player has full knowledge of his and his opponents' preferences over the outcomes, of pure strategies available to him and to his opponents, and that each player is aware that he and his opponents are all *rational* in that they all engage in a play of the game that is consistent with their respective preferences over the outcomes. In an ideal scenario then, players would seek out a play that produces an *optimal* outcome o , that is, an outcome that appears as the very next best one in the list of preferences of all the players. However, players typically end up seeking out a play that satisfies the constraints of their non-identical preferences over the outcomes. Such an outcome has the property that in the corresponding strategy profile (or the play of the game) each player's strategy is a *best reply* to all others, that is, any unilateral change of strategy in such a play of the game by any one player will produce an outcome of equal or less preference for that player than before. Such a play of a game is called a *Nash equilibrium* [4]. A Nash equilibrium that happens to correspond to an optimal outcome is called Pareto-optimal or simply an optimal Nash equilibrium.

Consider as an example the two player, two strategy non-cooperative game $\mathcal{G}$ with the set of outcomes $O = \{o_1, o_2, o_3, o_4\}$ and preferences of the players, labeled Player I and Player II, given by

$$\text{Player I : } o_2 \succ o_1 \succ o_4 \succ o_3 \quad (1.1)$$

$$\text{Player II : } o_3 \succ o_1 \succ o_4 \succ o_2 \quad (1.2)$$

with the symbol $\succ$ standing as short-hand for “more preferred than”. The set of pure strategies of each player is the two element set $\{D, H\}$ and the game is defined as

$$\mathcal{G}(D, D) = o_1, \quad \mathcal{G}(D, H) = o_3, \quad \mathcal{G}(H, D) = o_2, \quad \mathcal{G}(H, H) = o_4$$

Note from the players' preferences that the outcome o_1 is optimal, corresponding to the play (D, D) , because it best approximates the non-identical preferences of the players. However, the outcome o_4 is the unique Nash equilibrium in this game, corresponding to the strategy profile (H, H) , because it satisfies the constraint of the players' non-identical preferences over the outcomes. To see this, consider the game $\mathcal{G}$ in tabular format in Figure 1 from which it is easy to see that the player who unilaterally deviates from the strategy profile (H, H) ends up with an outcome that he prefers less than the outcome o_4 .

		Player II	
		D	H
Player I	D	o_1	o_3
	H	o_2	o_4

← Nash Equilibrium

Figure 1: Tabular form of the game $\mathcal{G}$.

1.1 Non-existent or Sub-optimal Nash Equilibria and Game Extensions

In the game $\mathcal{G}$ above, note that while an optimal outcome exists, it is not the one that manifests as the Nash equilibrium. This situation is considered to be an undesirable solution to the game. Worse game-theoretic situations can occur. For instance, the Matching Pennies game of Figure 2 entertains no optimal outcomes and no Nash equilibria! A fundamental problem in game theory is that of finding ways around such undesirable situations. One way this can be achieved is by simply changing the game. For instance, one can talk of changing the players' preferences in the game $\mathcal{G}$ or even changing the function $\mathcal{G}$ itself. However, changing a game can very naturally be viewed as a form of cheating. Indeed, a humorous lament betrays this assessment of changing a game as follows: "When I finally figured out answers to all of life's questions, they changed all the questions!" On a more practical note, one could argue that changing game as above is can be costly. For example, this cost of changing a game may be social, such as the one that Ghandi and the inhabitants of the Indian sub-continent incurred in their changing of the British Raj game.

Is there a way then to overcome undesirable game-theoretic situations where Nash equilibria are non-existent or sub-optimal, without changing the game? An affirmative answer can be found in the form of randomization via probability distributions. Randomization via probability distribution is a time-honored method of avoiding undesirable game-theoretic situations that has been practiced by players since time immemorial in the form of tossing coins, rolling die, or drawing straws, but was formally introduced in game theory by von Neumann [5].

2 Mixing the game

Randomization via probability distributions is introduced in a game formally as an exercise in the extension of the game. Bleiler provides an excellent treatment of this perspective in [1]. Taking the game $\mathcal{G}$ as an example, we extend its range to include probability distributions over the outcomes by identifying these outcomes with the corners of the simplex of

		Player II	
		Silent	Tell
Player I	Murder	(1, -1)	(-1, 1)
	Release	(-1, 1)	(1, -1)

Figure 2: Matching Pennies, a game with no Nash equilibrium.

probability distributions over four things, or the 3-simplex. Now probability distributions over the outcomes of $\mathcal{G}$ can be formed while at the same time any one of these original outcomes can be recovered by putting full probabilistic weight on that outcome. Formally, the set $O = \{o_1, o_2, o_3, o_4\}$ is identified with the set

$$\Delta_3 = \left\{ (p_1, p_2, p_3, p_4) \mid 0 \leq p_i \leq 1, \sum_{i=1}^4 p_i = 1 \right\}$$

where p_i is the probabilistic weight on the outcome o_i .

Next, a notion of non-identical preferences of the players over probability distributions is defined. A typical way to define such preferences is via the notion of *expectation*, constructed by assigning a numeric value, one per player, to each of the outcomes such that the assignment respects the preferences of the players over the outcomes. For a given probability distribution, expectation is defined to be

$$E(p_1, p_2, p_3, p_4) = p_1 a_1 + p_2 a_2 + p_3 a_3 + p_4 a_4 \in \mathbb{R} \quad (2.1)$$


with a_i being the numeric value of the outcome o_i . A player will now prefer one probability distribution P over another Q if $E(P) > E(Q)$. The *mixed* game M is defined next with domain equal to the Cartesian product of the sets of probability distributions over the pure strategies of the players. In symbols,

$$M : \Delta_1 \times \Delta_1 \rightarrow \Delta_3$$

where Δ_1 is the set of probability distributions over the pure strategies of each player and is referred to as the set of *mixed strategies* of the players. The mixed game M is defined as

$$M : ((p, 1 - p), (q, 1 - q)) \mapsto (pq, (1 - p)q, p(1 - q), (1 - p)(1 - q))$$

Figure 3 gives a pictorial representation of the construction of a mixed game for the game



$$M : ((p, 1-p), (q, 1-q)) \mapsto (pq, (1-p)q, p(1-q), (1-p)(1-q))$$

Figure 3: Construction of the mixed game M .

$\mathcal{G}$. We point out that the original game $\mathcal{G}$ can be recovered from the mixed game M by restricting the domain of M to $\{D, H\} \times \{D, H\}$. Hence, even though the mixed game is an entirely different game, because the original game is included within the mixed game and can be recovered if so desired, game theorists argue that the mixed game is one way around any undesirable game-theoretic results present in the original game *without changing the game*.

The existence of Nash equilibria in the mixed game was addressed by John Nash [4] who showed that a mixed game with a finite number of players is guaranteed to entertain at least one Nash equilibrium. This powerful result offers a way around the most undesirable situation possible in multi-player game non-cooperative theory, namely, a game without any any Nash equilibria. Moreover, it is sometimes the case that Nash equilibrium in the mixed game are near or fully optimal relative to the original game.

3 Gaming the Mixture

What can be said about the remains of the mathematical construction employed to produce the mixed game M when the underlying game $\mathcal{G}$ and its domain are removed from consideration? In this case, the function that remains, call it M' , maps directly from $\Delta_1 \times \Delta_1$ into Δ_4 and is no longer an extension of $\mathcal{G}$. The corners of each Δ_1 no longer correspond to mixed strategies of the players and the function $\mathcal{G}$ can no longer be recovered by introducing appropriate restrictions on the domain of M' . However, because the range of the game $\mathcal{G}$, together with players preferences defined over its elements, is still intact, included

inside Δ_4 , a notion of players' preferences over elements of Δ_4 as constructed in section 2 still holds. The new function M' can therefore be considered to be a multi-player non-cooperative game in which the set of outcomes is Δ_3 with players' preferences over these outcomes defined in terms of the players preferences over the corners of Δ_3 via expectation, and the “pure strategy” sets of each player equal Δ_1 :

$$M' : \Delta_1 \times \Delta_1 \rightarrow \Delta_4.$$

In other words, M' is the result of “gaming the mixture” or the application of multi-player non-cooperative game theory from section 1 to the stochastic function M' . Indeed, the function M' can appropriately be classified as a *stochastic game*.

This idea of casting a given function in a game-theoretic setting by starting with a game extension and then removing the underlying game from consideration is extended to functions used in quantum mechanics in section 5 below. To this end however, first a relevant discussion on extensions of games to include quantum mechanics appears below.


4 Quantizing the Game

Note that because M does not map onto Δ_3 , the image of the mixed game may not contain probability distributions that are optimal or near-optimal. Hence, the mixed game may not entertain Nash equilibria that are better those available originally. This is indeed the case with the game $\mathcal{G}$. In such persistent unsatisfactory situations, game-theorists seek other extensions of the original game.

Of special relevance here is the extension of a game, formally suggested by Meyer [2], to included higher order randomization via quantum superpositions followed by measurement. To this end, the outcomes of a game are identified with an orthogonal basis of the space of quantum superpositions. Mathematically, the space of quantum superpositions is a projective complex Hilbert space. For the game $\mathcal{G}$, a four dimensional projective Hilbert space $\mathbf{H}_4$ is required with the four outcomes $\{o_1, o_2, o_3, o_4\}$ identified with an orthogonal basis $B = \{b_1, b_2, b_3, b_4\}$ of $\mathbf{H}_4$. Now quantum superpositions of the outcomes of $\mathcal{G}$ can be formed. Formally, the set $O = \{o_1, o_2, o_3, o_4\}$ is identified with the set

$$\mathbf{H}_4 = \left\{ c_1 b_1 + c_2 b_2 + c_3 b_3 + c_4 b_4 = (c_1, c_2, c_3, c_4) \mid \sum_{i=1}^4 |c_i|^2 = 1 \right\}$$

where c_i is the projective complex valued weight on o_i and $|c_i|^2$ is the square of the norm of



$$\Theta: (\alpha_1 D + \alpha_2 H, \beta_1 D + \beta_2 H) \mapsto (\gamma_1, \gamma_2, \gamma_3, \gamma_4)$$

Figure 4: Construction of quantized games Θ .

the complex number c_i . Measurement, denoted here as *meas*, of a quantum superposition q produces a probability distribution in Δ_3 from which the expectation of the outcomes of $\mathcal{G}$ to the players can be computed and the optimality of a quantum superposition can be defined. Note that anyone of the outcomes of the game $\mathcal{G}$ can be recovered by putting full quantum superpositional weight on the basis element corresponding to that outcome. A *quantized game* Θ is defined next, typically as a length preserving unitary function from the Cartesian product of the sets of quantum superpositions of the strategies of the players into $\mathbf{H}_4$. In symbols,

$$\Theta: \mathbf{H}_2 \times \mathbf{H}_2 \rightarrow \mathbf{H}_4$$

where $\mathbf{H}_2$ denotes the set of quantum superpositions of the strategies of the players, referred to in the literature as *quantum strategies* of the players. However, because a quantized game is an extension of the original game $\mathcal{G}$ in a fashion analogous to the mixed game of section 2, we propose that it is more appropriate to refer to $\mathbf{H}_2$ as the set of a player's *quantized strategies*. We also point out that the term quantized game in fact refers to an entire family of games, in stark contrast to the term mixed game which refers to a specific function. Depending on the exact nature of Θ , when followed by measurement, the image of this composite map $\text{meas} \circ \Theta$ can be larger than the image of mixed game and may contain optimal or near-optimal probability distributions. A pictorial representation of a game quantization is given in Figure 4.

Quantization of games with two players, each having two pure strategies have been

fairly extensively studied, with a recent survey of this subject appearing in [6]. One quantization that underpins most studies of two player, two strategy games is the one proposed by Eisert, Wilkens, and Lewenstein (EWL) [3]. It can be shown that the EWL quantization is the specific family of functions explicitly defined as

$$\Theta : (\alpha_1 D + \alpha_2 H, \beta_1 D + \beta_2 H) \rightarrow (\gamma_1, \gamma_2, \gamma_3, \gamma_4)$$

where

$$\begin{aligned}\gamma_1 &= (\alpha_1 \beta_1 + \alpha_2 \beta_2) + (\bar{\alpha}_1 \bar{\beta}_1 + \bar{\alpha}_2 \bar{\beta}_2) \\ \gamma_2 &= -\eta(-\alpha_1 \bar{\beta}_2 + \alpha_2 \bar{\beta}_1) + \bar{\eta}(-\bar{\alpha}_2 \beta_1 + \bar{\alpha}_1 \beta_2) \\ \gamma_3 &= \bar{\eta}(-\alpha_1 \bar{\beta}_2 + \alpha_2 \bar{\beta}_1) - \eta(-\bar{\alpha}_2 \beta_1 + \bar{\alpha}_1 \beta_2) \\ \gamma_4 &= -i(\alpha_1 \beta_1 + \alpha_2 \beta_2) + i(\bar{\alpha}_2 \bar{\beta}_2 + \bar{\alpha}_1 \bar{\beta}_1) \\ \eta &= e^{\frac{i\pi}{4}} = \frac{1}{\sqrt{2}}(1 + i),\end{aligned}$$

and focuses on a particular variation of the game $\mathcal{G}$ with specific numeric values replacing the outcomes o_i . This is the popular game known as Prisoner's Dilemma. The authors of EWL show that a Nash equilibrium with an expectation equal to the optimal outcome in the original game $\mathcal{G}$ manifests in the quantized game for a play of the game consisting of a certain sub-class of quantized strategies. However, when plays of the game consisting of the most general class of quantized strategies are considered, the only Nash equilibrium that manifests is one with expectation equal to that of the sub-optimal Nash equilibrium in the original game. Indeed, Meyer presents an argument in [2] where he shows that unlike a mixed game, a quantized game need not entertain any Nash equilibria at all.

5 Gaming the Quantum

Just as removing the underlying game $\mathcal{G}$ and its domain from consideration in the construction of the mixed game M in section 2 leaves the function M' that can be viewed as an example of applying multi-player non-cooperative game theory to stochastic functions, removing the game $\mathcal{G}$ from consideration in the construction of a quantized game Θ leaves behind a function Θ' that can be viewed as an example of an application of multi-player, non-cooperative game theory to quantum mechanics or “gaming the quantum”. Indeed, because the function Θ' is a quantum physical function (not necessarily a unitary operator, but *any* function in general) by virtue of mapping into the joint state space of quantum me-




Figure 5: Distance between a quantum superposition Q and elements of b_i of an orthogonal basis of $\mathbf{H}_4$.

chanical objects, it truly deserves to be called a *quantum game*. In other words, a quantum game is any function mapping into a projective complex Hilbert space provided a notion of preferences, one per player, is defined over quantum superpositions. The domain of a quantum game can now be correctly referred to as the set of *quantum strategies* of the players. Indeed, in a general setting, the set of quantum strategies can be any set and need not be restricted to $\mathbf{H}_2$.

While the notion of players' preferences over quantum superpositions in the outcome space $\mathbf{H}_4$ can conceivably be defined in different ways, here, we define players' preferences over these quantum superpositions via preferences over the elements of an orthogonal basis of $\mathbf{H}_4$ in a particular way. Define the preferences of the players over the elements of orthogonal basis $B = \{b_1, b_2, b_3, b_4\}$ of $\mathbf{H}_4$ to be preferences over the elements of an orthogonal basis of $\mathbf{H}_4$ in a particular way. Define the preferences of the players over the elements of orthogonal basis $B = \{b_1, b_2, b_3, b_4\}$ of $\mathbf{H}_4$ to be

$$\text{Player I} : b_2 \succ b_1 \equiv b_4 \equiv b_3$$

$$\text{Player II} : b_3 \succ b_1 \equiv b_4 \equiv b_2$$

where the symbol " $\equiv$ " represents a players indifference between the basis elements surrounding the symbol. In other words, Player I prefers the basis element b_2 the most and Player II prefers the basis element b_3 the most. Next, note that the Hilbert space $\mathbf{H}_4$ natu-

rally supports a notion of distance (metric, in fact) between any two of its elements via the inner-product defined on it. More precisely, for $x_1, x_2 \in \mathbf{H}_4$, the distance between these two elements is given by the angle

$$\text{dist}(x_1, x_2) = \theta_{(x_1, x_2)} = \cos^{-1} (| \langle x_1, x_2 \rangle |)$$

where $\langle x_1, x_2 \rangle$ is the inner-product of x_1 and x_2 , $| \langle x_1, x_2 \rangle |$ represents its length or norm, and $\theta \in [0, \frac{\pi}{2}]$. This notion of distance is also referred to as the *fidelity* of two quantum superpositions.

Player I will now prefer one quantum superposition q over another p if q is closer to b_2 than p is, that is

$$\text{Player I : } q \succ p \Leftrightarrow \text{dist}(q, b_2) < \text{dist}(p, b_2).$$

Similarly, Player II will prefer one quantum superposition q over another r if q is closer to b_3 than r is:

$$\text{Player II : } q \succ r \Leftrightarrow \text{dist}(q, b_3) < \text{dist}(r, b_3)$$

Now the notion of Nash equilibrium can be characterized as a play of a quantum game that satisfies the constraints of the players' preferences via this distance notion. Let Q be a quantum superposition corresponding to a play (x^*, y^*) of a quantum game Γ , that is

$$\Gamma(x^*, y^*) = Q$$

The quantum superposition Q will be a Nash equilibrium outcome if unilateral deviation on part of any one player from the corresponding play (x^*, y^*) of Θ will produce a quantum superposition of equal or less preference for *that* player than Q . Therefore, if Player I deviates from his quantum strategy x^* and instead employs the quantum strategy x , then

$$\Gamma(x, y^*) = Q_x \quad \text{with} \quad \text{dist}(Q_x, b_2) \geq \text{dist}(Q, b_2) \quad (5.1)$$

Also, if Player II deviates from his quantum strategy y^* and instead employs the quantum strategy y , then

$$\Gamma(x^*, y) = Q_y \quad \text{with} \quad \text{dist}(Q_y, b_3) \geq \text{dist}(Q, b_3) \quad (5.2)$$

Note that equations (5.1) and (5.2) describe a simultaneous best approximation problem in the Hilbert space $\mathbf{H}_4$. This establishes the main result:

A quantum strategy pair (s^*, t^*) in a two player quantum game Γ is a Nash equilibrium if $\Gamma(s^*, t^*) = Q \in \mathbf{H}_4$ simultaneously best approximates the most preferred outcomes of the players.

The general case with finitely many players follows straight forwardly.

The theory of Hilbert space shows that for a given sub-Hilbert $\mathcal{S}$ space of a Hilbert space $\mathcal{H}$, one can always find a unique element $s \in \mathcal{S}$ that minimizes the distance between elements of $\mathcal{S}$ and any fixed $h \in \mathcal{H}$. In other words, s best approximates h . This means that a *necessary* condition for the existence of Nash equilibrium in a quantum game is for the image of the quantum game to be a sub-Hilbert space of the $\mathbf{H}_4$. On the other hand, equations (5.1) and (5.2) constitute a *sufficient* condition for the existence of Nash equilibrium in a quantum game.

6 Conclusion

We envision several potential future directions in the area of quantum games. For one, players' preferences given in section 5 above are defined within a quantum computational and algorithmic context where exactly one outcome, corresponding to a particular calculation or searched item, is the correct and therefore most desired outcome of one player. All others are less preferable. On the other hand, another player (or even players) most prefers at least one of the latter. The equilibrium behavior of quantum circuits can potentially be studied from this game-theoretic perspective. Indeed, other preferences over quantum superpositions are conceivable, and an entire separate project can be devoted to the study preferences that produce insightful results.

Further generalization is another possible future direction. One would start with the study of the class of functions into $\mathbf{H}_4$, or indeed into $\mathbf{H}_n$ for any $n \in \mathbb{N}$, that would satisfy the necessary and sufficient conditions for the existence of Nash equilibria. Generalizing further would allow a game-theoretic study of a broader class of functions, culminating with the positive operator valued measurement mapping into infinite dimensional vector spaces with a continuum of basis elements. Such studies can potentially produce insightful results in the engineering of control of quantum systems those corresponding to quantum circuits and algorithms.

7 Acknowledgments

The authors gratefully acknowledge useful discussions with Steven Bleiler and Joel Lucero-Bryan.

References

- [1] S. A. Bleiler, *A Formalism for Quantum Games I - Quantizing Mixtures*, Portland State University, preprint at <http://arxiv.org/abs/0808.1389>, 2008. 3
- [2] D. Meyer, *Quantum Strategies*, Physical Review Letters, Volume 82, pages 1052-1055, 1999. 6, 8
- [3] J. Eisert and M. Wilkens and M. Lewenstein, *Quantum Games and Quantum Strategies*, Physical Review Letters, Volume 83, pages 3077-3080, 1999. 8
- [4] J. F. Nash *Equilibrium Points in n -Person Games*, Proceedings of the National Academy of Sciences of the United States of America, Issue 36, pages 48-49, 1950. 2, 5
- [5] J. V. Neumann, *Theory of Games and Economic Behavior*, Princeton University Press, 1944. 3
- [6] S. Lansburg, *Quantum Game Theory*, Wiley Encyclopedia of Operations Research and Management Science, 2011. 8